



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities.

ALABAMA—*Mobile*.—Month of January, 1897. Estimated population, 31,076. Total deaths, 98, including phthisis pulmonalis, 12; enteric fever, 1, and croup, 1.

CALIFORNIA—*Los Angeles*.—Month of January, 1897. Estimated population, 100,000. Total deaths, 131, including phthisis pulmonalis, 37; enteric fever, 2, and diphtheria, 2.

Oakland.—Month of January, 1897. Estimated population, 36,000. Total deaths, 73, including phthisis pulmonalis, 12; enteric fever, 1, and measles, 1.

Sacramento.—Estimated population, 30,000. Total deaths, 48, including phthisis pulmonalis, 9, and diphtheria, 1.

COLORADO—*Denver*.—Month of January, 1897. Estimated population, 106,713. Total deaths, 166, including phthisis pulmonalis, 47; enteric fever, 2; scarlet fever, 1; diphtheria, 8; whooping cough, 1, and measles, 3.

CONNECTICUT—*Hartford*.—Month of January, 1897. Estimated population, 72,000. Total deaths, 103, including phthisis pulmonalis, 9; enteric fever, 1; scarlet fever, 1; diphtheria, 4, and whooping cough, 2.

FLORIDA.—Month of November, 1896. Reports to the State board of health from 45 counties, including the cities of Jacksonville, Key West, and Pensacola, having an aggregate population of 391,418, show a total of 249 deaths, including enteric fever, 7; phthisis pulmonalis, 34; croup, 1, and diphtheria, 1.

Month of December, 1896. Total deaths, 296, including smallpox, 1; phthisis pulmonalis, 38; enteric fever, 7; scarlet fever, 1; diphtheria, 3, and croup, 3.

LOUISIANA—*Shreveport*.—Month of January, 1897. Estimated population, 15,000. Total deaths, 33, including 6 from phthisis pulmonalis.

MASSACHUSETTS—*Brockton*.—Month of January, 1897. Estimated population, 33,165. Total deaths, 51, including phthisis pulmonalis, 9; diphtheria and croup, 5, and measles, 1.

Newton.—Month of January, 1897. Total deaths, 38, including phthisis pulmonalis, 3, and enteric fever, 1.

MICHIGAN.—Week ended February 6, 1897. Reports to the State board of health, Lansing, from 57 observers, indicate that measles, erysipelas, and inflammation of kidney increased and consumption decreased in era of prevalence. Phthisis pulmonalis was reported present during the week at 161 places, measles at 51, diphtheria at 42, scarlet fever at 26, enteric fever at 16, and whooping cough at 15 places.

Grand Rapids.—Month of January, 1897. Estimated population, 90,000. Total deaths, 84, including phthisis pulmonalis, 11; enteric fever, 3; croup, 1; diphtheria, 1, and scarlet fever, 1.

MINNESOTA—*Winona*.—Month of January, 1897. Estimated popula-

tion, 22,000. Total deaths, 36, including phthisis pulmonalis, 3; croup, 1, and whooping cough, 1.

NEW HAMPSHIRE—*Concord*.—Month of January, 1897. Estimated population, 19,000. Total deaths, 36, including 1 from diphtheria.

Manchester.—Month of January, 1897. Estimated population, 56,000. Total deaths, 128, including phthisis pulmonalis, 8; diphtheria, 7; croup, 2, and whooping cough, 2.

PENNSYLVANIA—*Pottsville*.—Two weeks ended February 11, 1897. Estimated population, 14,000. Total deaths, 16, including 1 from diphtheria.

TENNESSEE—*Knoxville*.—Month of January, 1897. Estimated population, 45,000; white, 35,000; colored, 10,000. Total deaths, 58—white, 43, colored, 15—including phthisis pulmonalis, 10; enteric fever, 2, and measles, 2.

Memphis.—Month of January, 1897. Estimated population—white, 28,573; colored, 27,561; total, 56,134. Deaths—white, 57; colored, 57; total, 114, including phthisis pulmonalis, 24; enteric fever, 2, and diphtheria, 1.

WASHINGTON—*Seattle*.—Month of January, 1897. Estimated population, 60,000. Total deaths, 32, including phthisis pulmonalis, 6, and enteric fever, 2.